

## § 522.2680

(1) *Amount.* 0.2 to 0.3 milligram per kilogram of body weight.

(2) *Indications for use.* As an antagonist to xylazine sedation in free ranging or confined members of the family Cervidae (deer and elk).

(3) *Limitations.* For intravenous use only. Do not use in domestic food-producing animals. Do not use for 30 days before or during hunting season. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 8543, Feb. 16, 1993, as amended at 60 FR 57832, Nov. 22, 1995]

## § 522.2680 Zeranor.

(a) *Specifications.* Each pellet contains 12, 18, or 20 milligrams (mg) zeranol.

(b) *Sponsor.* See 000061 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.760 of this chapter.

(d) *Conditions of use*—(1) *Beef cattle*—(i) *Amount.* 36 mg zeranol (one implant consisting of 3 pellets, each pellet containing 12 mg zeranol) per implant dose.

(ii) *Indications for use*—(A) For increased rate of weight gain and improved feed conversion in weaned beef calves, growing beef cattle, feedlot steers, and feedlot heifers.

(B) For increased rate of weight gain in suckling calves.

(iii) *Limitations.* Implant subcutaneously in ear only. Do not use in bulls intended for reproduction or in dairy animals. Do not use before 1 month of age or after weaning in heifers intended for reproduction.

(2) *Feedlot lambs*—(i) *Amount.* 12 mg zeranol (one implant consisting of 1 pellet containing 12 mg zeranol) per implant dose.

(ii) *Indications for use.* For increased rate of weight gain and improved feed conversion.

(iii) *Limitations.* Implant subcutaneously in ear only. Do not use in breeding animals. Do not implant animals within 40 days of slaughter.

(3) *Steers fed in confinement for slaughter*—(i) *Amount.* 72 mg zeranol (one implant consisting of 6 pellets, each pellet containing 12 mg zeranol) per implant dose.

## 21 CFR Ch. I (4–1–02 Edition)

(ii) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(iii) *Limitations.* Implant subcutaneously in ear only.

(4) *Pasture cattle (slaughter, stocker, feeder steers, and heifers)*—(i) *Amount.* 138 mg zeranol (one implant consisting of 7 pellets, each of 6 pellets containing 20 mg zeranol and a seventh pellet containing 18 mg zeranol) per implant dose.

(ii) *Indications for use.* For increased rate of weight gain.

(iii) *Limitations.* Implant subcutaneously in ear only.

[59 FR 19639, Apr. 25, 1994; 60 FR 26360, May 17, 1995, as amended at 62 FR 61625, Nov. 19, 1997; 64 FR 46840, Aug. 27, 1999; 67 FR 6867, Feb. 14, 2002]

## PART 524—OPHTHALMIC AND TOPICAL DOSAGE FORM NEW ANIMAL DRUGS

Sec.

524.86 Amitraz liquid.

524.154 Bacitracin or bacitracin zinc-neomycin sulfate-polymyxin B sulfate ophthalmic ointment.

524.155 Bacitracin zinc-polymyxin B sulfate-neomycin sulfate-hydrocortisone or hydrocortisone acetate ophthalmic ointment.

524.390 Chloramphenicol ophthalmic and topical dosage forms.

524.390a Chloramphenicol ophthalmic ointment.

524.390b Chloramphenicol ophthalmic solution.

524.390d Chloramphenicol-prednisolone ophthalmic ointment.

524.402 Chlorhexidine ointment.

524.450 Clotrimazole cream.

524.463 Copper naphthenate solution.

524.520 Cuprimyxin cream.

524.575 Cyclosporine ophthalmic ointment.

524.660 Dimethyl sulfoxide ophthalmic and topical dosage forms.

524.660a Dimethyl sulfoxide solution.

524.660b Dimethyl sulfoxide gel.

524.770 Doramectin.

524.802 Enrofloxacin, silver sulfadiazine emulsion.

524.814 Eprinomectin.

524.900 Famphur.

524.920 Fenthion.

524.960 Flumethasone, neomycin sulfate, and polymyxin B sulfate ophthalmic solutions.

524.981 Fluocinolone acetonide ophthalmic and topical dosage forms.

524.981a Fluocinolone acetonide cream.